

LESSON 12

- **MODIFIERS AND MODIFIED EXPRESSIONS**
 - Common Modifiers
 - Binomial Coefficient
 - Modified Expressions and Superscripts / Subscripts
 - Modified Signs of Comparison
 - Expressions with More Than One Modifier

Format

- Formal Proof

Postscript: Vector Notation Notes

Answers to Practice Material

LESSON PREVIEW

Another type of mathematical notation with vertical components is studied. This lesson also applies the "mathematical statement" format to formal proofs.

Some of the print examples are enlarged in order to more clearly show the modifiers on the printed page. You may also wish to use a magnifier to remove any uncertainty.

MODIFIERS AND MODIFIED EXPRESSIONS

[NC Rule 15]

Some mathematical notation has a vertical aspect that challenges linear braille notation. By using modified expression indicators, the transcriber can relay the material in a compact manner, and the reader can construct the symbols into a meaningful expression.

12.1 Definition

A modifier is a symbol or a combination of symbols occurring *directly over* or *directly under* its related symbol or expression. Here are some typical examples.

$\overset{?}{=}$ a question mark over an equals sign

$\underline{3.15}$ an underlined digit

$1.\overline{37}$ a line over two numerals (signifying a repeating decimal)

\overleftrightarrow{AB} an arrow over two letters (signifying a line)

\hat{k} a caret over a letter

12.2 Construction of Simple Modified Expressions – The Five-Step Rule

An expression modified using the five-step method is initiated and terminated with special indicators.

$\ddot{\cdot}$	Multipurpose Indicator
$\ddot{\cdot}$	Termination Indicator

The position of the modifier (above or below the expression) is also shown with the use of an indicator.

$\overset{\cdot}{\cdot}$	Directly-Over Indicator
$\underset{\cdot}{\cdot}$	Directly-Under Indicator

The process of constructing a modified expression is known as "The Five-Step Rule".

- (1) The *multipurpose indicator* is placed immediately before the expression to be modified.
- (2) The *expression* to be modified is written second.

- (3) The *directly-over indicator* or the *directly-under indicator* is written third to show the position of the modifier.
- (4) The *modifying symbol* is written fourth.
- (5) The *termination indicator* is written last to show the end of the modification.

Notice that the multipurpose indicator, dot 5, signals the beginning of the modified expression and the termination indicator signals the completion of the modified expression.

Prototype for an expression with a modifier printed directly above it: $\cdot \underline{\quad} \cdot \underline{\quad} \cdot$

Prototype for an expression with a modifier printed directly below it: $\cdot \underline{\quad} \cdot \underline{\quad} \cdot$

All components of an expression modified according to the five-step rule should not be divided between lines. If that is not possible, strategies will be presented in Lesson 15.

Common Modifiers

The most commonly used modifiers are presented in this lesson. You have seen many of these symbols in other contexts in previous lessons. Symbols of the code not shown here may also be used as modifiers.

12.3 Arrows as Modifiers

Any of the arrows of the Nemeth Code may be a modifier. Those used in this section are shown below.

$\cdot \cdot$	Arrow barbed at right, contracted form	\rightarrow
$\cdot \cdot \cdot \cdot \cdot$	Arrow barbed at both ends	\leftrightarrow
$\cdot \cdot \cdot \cdot \cdot$	Arrow shaft with hollow dot at right	$\rightarrow\circ$
$\cdot \cdot \cdot \cdot \cdot$	Upper barb only, right pointing	\rightarrow
$\cdot \cdot \cdot \cdot \cdot$	Arrow barbed at right, dashed shaft	\dashrightarrow

- 12.3.1 **Right-Pointing Arrow.** A right-pointing arrow in regular type with a full barb and single shaft of ordinary length is transcribed in its contracted form when used as a modifier above or below a math expression. The shaft length in print is determined by the width of the expression it modifies. If the modified expression is wide, the arrow shaft will be long in print, but this is not considered to be a "longer than ordinary" arrow shaft in this context.

Example 12-4

Use arrow math to add $36 + 23$.

$36 \xrightarrow{+20} 56 \xrightarrow{+3} 59 \quad 36 + 23 = 59$

Instructions: Review arrow construction in Lesson 9. *Reminder:* In braille, identifiers for nonspatial itemized material must all begin on a new line in cell 1.

In the sentence at the end of this practice, assume that all vectors in the document are shown using that particular arrow notation. Show the proper way to omit the vector arrows in the transcription. Include the required transcriber's note after the topic heading.

PRACTICE 12A

Here are two modified arrows: $x \xrightarrow{g} y \xrightarrow{f} z$

Arrows as Modifiers

1. \overleftarrow{F}
2. \overleftarrow{AB}
3. \overleftarrow{CD}
4. $\overleftrightarrow{OB} \cup \overleftrightarrow{OC}$
5. $\overset{\circ}{\overleftarrow{EF}}$
6. $\overleftarrow{\overleftarrow{T}}$
7. $\overleftrightarrow{XZ} \parallel \overleftrightarrow{RS}$
8. $\overleftrightarrow{AB} + \overleftrightarrow{CD}$

Vector Addition

\mathbb{R} equals \overrightarrow{OP} equals \overrightarrow{OM} plus \overrightarrow{MC} plus \overrightarrow{CP} .

Instructions: Treat the three examples of unit vectors in problem #2 as displayed mathematical material.

PRACTICE 12B

Carets and Bars as Modifiers

1. Unit vectors can be denoted with normal vector notation, \mathbf{u} or \vec{u} , or with standard unit vector notation, $\hat{\mathbf{u}}$, spoken "u-hat".
2. Unit vectors in various coordinate systems use Greek and English letters.

Cartesian coordinate system: $\hat{\mathbf{x}}, \hat{\mathbf{y}}, \hat{\mathbf{z}}$

Cylindrical coordinate system: $\hat{\rho}, \hat{\phi}, \hat{\mathbf{z}}$

Spherical coordinate system: $\hat{\mathbf{r}}, \hat{\theta}, \hat{\phi}$

3. $1.142857\overline{142857}$

4. $\frac{7}{15} = .4\overline{6}$

5. $2 \cdot 3 = \overline{2} \cdot 3 = \overline{2 \cdot 3}$

6. $\overline{PQ}, \overline{x'}, \overline{R''S''}$

7. $\overline{s}, \overline{\alpha}, \overline{m'}$

8. $m\overline{BC} = a$

9. $\overline{C} = 100 \times 1000$

10. $F = 2\pi\overline{r}l$


11. $P(\overline{a + bi}) = \overline{0} = 0$

12. $\hat{x}_i = 0.5(\overline{x}_i + \underline{x}_i)$

13. $3.141\underline{59}$


Example 12-12

Expressed as a fraction, $0.909\dot{0}\dot{9} = \frac{1}{11}$.




In print, a single dot is shown over each of the last two digits (0 and 9).

When one or more dots occur over or under a single letter or numeral, the symbol for the dot is used as many times as necessary to conform with the print text.




Example 12-13

Prove $\ddot{x} = \frac{d^2x}{dt^2}$



12.6.3 **Horizontal Grouping Sign.** When a horizontal grouping sign occurs over or under a mathematical expression, it is either a part of the expression or is pointing to a label.

- a. **As a Pointer.** When the grouping sign points to a label or to explanatory text, it must be drawn as a tactile graphic. Refer to *Guidelines and Standards for Tactile Graphics* for drawing techniques.




Example 12-14


To simplify the polynomial, combine like terms.

$$(2x + 3)(x - 5) = 2x^2 - 10x + 3x - 15$$


$$= 2x^2 - 7x - 15$$



- b. **As a Modifier.** When a horizontal grouping sign is part of the math expression, it is regarded as a modifier. Although it is preferable to draw the symbol, the transcriber can represent the construction as shown below, following the five-step rule of modified expressions. The left grouping symbol is transcribed when the modifier is directly over the expression, and the right grouping symbol is transcribed when the modifier is directly under the expression.



- 12.6.4 **Question Mark.** When the question mark is not functioning as a sign of omission, the punctuation mark is transcribed. In mathematical context, a punctuation indicator precedes the question mark to prevent it from being misread as the numeral 8.



An equals sign is the expression being modified. It is modified directly over with a question mark.

Greek capital letters *Sigma* and *Pi* are often encountered in this notation. Sigma is used in summation notation and Pi is used in product notation.

⠠⠠⠠⠠⠠	Sigma	Σ	(capital)
⠠⠠⠠⠠⠠	Pi	Π	(capital)

$$\Rightarrow \sum_{i < j} \quad \text{⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠}$$

Sigma is modified directly under with the inequality i < j.

$$\Rightarrow \prod_{k=1} \quad \text{⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠}$$

Pi is modified directly under with the equality k = 1.

12.7.1 Binomial Coefficient. A binomial coefficient is written as two expressions, one atop the other, enclosed in parentheses. For example, *n choose q* is written like this:

$$\binom{n}{q}$$

The binomial coefficient does not follow the five-step rule for modified expressions. The opening parenthesis is followed by the upper expression; the directly-under indicator is placed next, followed by the lower expression. The closing parenthesis ends the expression.

⠠⠠⠠⠠	Directly-under indicator
------	--------------------------

$$\Rightarrow \binom{n}{q} \quad \text{⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠}$$

$$\Rightarrow \binom{n-1}{k-1} \quad \text{⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠⠠}$$

Example 12-16

The **recursive formula** states that $\binom{n}{0} = \binom{n}{n} = 1$ for all integers $n, k : 1 \leq k \leq n - 1$.


$$\begin{array}{cccccccccccc} \text{⠠⠠⠠⠠} & \text{⠠⠠⠠⠠⠠⠠} & \text{⠠⠠⠠⠠⠠⠠⠠⠠} & \text{⠠⠠⠠⠠⠠⠠⠠⠠⠠} & \text{⠠⠠⠠⠠⠠⠠⠠⠠⠠} & \text{⠠⠠⠠⠠⠠⠠⠠} & \text{⠠⠠⠠⠠⠠⠠} & \text{⠠⠠⠠⠠} & \text{⠠⠠⠠⠠} & \text{⠠⠠⠠⠠} & \text{⠠⠠⠠⠠} & \text{⠠⠠⠠⠠} & \text{⠠⠠⠠⠠} & \text{⠠⠠⠠⠠} \\ \text{⠠⠠⠠⠠} & \text{⠠⠠⠠⠠⠠⠠} & \text{⠠⠠⠠⠠} & \text{⠠⠠⠠⠠⠠⠠} & \text{⠠⠠⠠⠠} & \text{⠠⠠⠠⠠} & \text{⠠⠠⠠⠠} & \text{⠠⠠⠠⠠} & \text{⠠⠠⠠⠠} & \text{⠠⠠⠠⠠} & \text{⠠⠠⠠⠠} & \text{⠠⠠⠠⠠} & \text{⠠⠠⠠⠠} & \text{⠠⠠⠠⠠} \\ \text{⠠⠠⠠⠠} & \text{⠠⠠⠠⠠} & \text{⠠⠠⠠⠠} & \text{⠠⠠⠠⠠} & \text{⠠⠠⠠⠠} & \text{⠠⠠⠠⠠} & \text{⠠⠠⠠⠠} & \text{⠠⠠⠠⠠} & \text{⠠⠠⠠⠠} & \text{⠠⠠⠠⠠} & \text{⠠⠠⠠⠠} & \text{⠠⠠⠠⠠} & \text{⠠⠠⠠⠠} & \text{⠠⠠⠠⠠} \end{array}$$

Reminder from Lesson 5: A colon that means "such that" is transcribed unspaced from the letter it follows, and is preceded by a punctuation indicator. This spacing is applied regardless of the spacing shown in print. Spacing after the colon follows print.

Example 12-17

Use this formula to calculate the binomial coefficient.

$$\binom{n}{k} = \frac{n!}{k! \cdot (n-k)!}$$



PRACTICE 12C


1. In Figure 7.3, if $\widehat{AB} = \widehat{CD}$ in circle O, then $\angle AOB = \angle BOC$.
2. $x \stackrel{?}{=} y$ means "does x equal y ?"
3. $\sum_{d|n}$ (where $d|n$ means " d divides n ").
4. $\binom{t}{p} = R_t^p$
5. $\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$ for all integers $n \geq 0$.
6. Does (\tilde{x}, \tilde{y}) mean \tilde{x} and \tilde{y} ?
7. More modified expressions:
 - a. $.249\grave{9}$
 - b. $2.431\dot{3}\hat{1}$
 - c. $\acute{x}\grave{y} - \grave{y}\acute{x}$
 - d. $\grave{a} + \grave{\grave{a}} = ?$
 - e. $f \rightarrow \tilde{f}$
 - f. $\widehat{x+y}$

12.8 Spacing with Modified Expressions

The spacing before and after an entire modified expression is subject to the spacing rules for the symbols preceding or following it.


Example 12-18

Prove that $\overline{OP} + \overline{QR} = \overline{OR}$.



There is no space before or after the operation (plus) sign. There is a space before and after the comparison (equals) sign.

When the modifier is wider than the modified symbol, the print copy will insert extra space to clarify what exactly is being modified. In the next example, only the Sigma is modified, not the letter a . The space between the Sigma and the a clarifies the extent of the modifier, $i < j$. In braille, however, indicators define the extent of a modifier. There is no need for the space in braille.



*Sigma is modified directly under with the inequality $i < j$.
The termination indicator signals the completion of the modifier.*

PRACTICE 12D

Spacing with Modified Expressions

A. The probability of the event A, written $P(A)$, is defined as

$$P(A) = \sum_{\wedge} f(x)$$

where $\sum_{\wedge} f(x)$ means sum $f(x)$ over those values x_i that are in A.

B. $\prod_{\alpha \in A} A_{\alpha}$

C. $\prod_{i > j} (x_i - x_j)$

D. $\overline{7} + 2 \stackrel{?}{=} \overline{7 + 2}$

$$\gg \overline{OD^2} + \overline{OP^2}$$



The first dot 5 begins the first modified expression. The second dot 5 is a baseline indicator following the first superscript "2". Similarly, the third dot 5 begins the second modified expression and the fourth dot 5 is a baseline indicator following the second superscript.

$$\gg \overline{x_1} \quad \dots$$

You may wish to review Section 6.11 regarding nonuse of the subscript indicator.

- a. **Binomial Coefficient.** Although a binomial coefficient is not technically a modified expression, notice how this rule applies.

$$\gg \binom{a_x}{b_y} \quad \dots$$

The baseline indicator precedes the "directly under" indicator. This keeps that indicator on the same level of writing as the letters "a" and "b".

PRACTICE 12E

Superscripts and Subscripts

- A) $\overline{AB}^2 + \overline{BC}^2$
- B) $\overline{A} = [\overline{a_i}]$
- C) $\sqrt{\overline{\dot{x}^2} + \overline{\dot{y}^2}}$
- D) \underline{Z}°
- E) If $\overline{a_1} = 72$, find $\overline{a_7}$.
- F) Draw $\overline{P_1P_2}$ if P_1 is the point (1, 3) and P_2 is the point (2, -1).
- G) $(\overline{3^{-1}}) \in P$
- H) $\overline{x_1} + \overline{y_1}$

Modified Signs of Comparison

12.11 Definition

A modified sign of comparison consists of a simple sign of comparison, such as the equals sign or the tilde, modified by a caret, dot, triangle, question mark, vertical bar, or any symbol except another sign of comparison.

When a simple sign of comparison occurs above or below another simple sign of comparison the combination is transcribed as a sign of comparison compounded vertically. See Section 5.8 for a review of that construction. Note that many of those signs are printed with a single horizontal line "bar over" or "bar under". For example, "bar over greater than, inclusion with bar under, bar over single tilde, logical sum with bar under," etc. These signs are not to be misinterpreted as a horizontal bar modifying a sign of comparison.

12.12 Transcription

A modified sign of comparison as defined above is transcribed in accordance with the five-step rule for modified expressions.

In addition to the caret and inverted caret seen earlier in this lesson, you may also encounter a left- or right-pointing caret in a modified sign of comparison. Do not confuse these two symbols with the "less than" and "greater than" comparison signs. Ask an expert if context does not clarify the identity of this symbol.

⠠⠠⠠⠠⠠⠠	Left-Pointing Caret	<
⠠⠠⠠⠠⠠⠠	Right-Pointing Caret	>

The following list contains the modified equals signs most commonly used.

Modified Equals Sign		
⠠⠠⠠⠠⠠⠠⠠⠠	Caret Over Equals Sign	≐
⠠⠠⠠⠠⠠⠠⠠⠠	Caret Under Equals Sign	≑
	("is projective to")	^
⠠⠠⠠⠠⠠⠠⠠⠠	Inverted Caret Over Equals Sign	≒
⠠⠠⠠⠠⠠⠠⠠⠠	Left-Pointing Caret Over Equals Sign	≓
⠠⠠⠠⠠⠠⠠⠠⠠	Right-Pointing Caret Over Equals Sign	≔

$\dot{=}$	Dot Over Equals Sign ("is approximately equal to")	\approx
$\underset{\cdot}{=}$	Dot Over and Dot Under Equals Sign	\cong
$\overset{\cdot\cdot}{=}\underset{\cdot\cdot}{=}$	Two Dots Over and Two Dots Under Equals Sign	\equiv
$\circ=$	Hollow Dot Over Equals Sign ("is equal in degrees to")	$\overset{\circ}{=}$
$\triangle=$	Equilateral Triangle Over Equals Sign	\triangleq
$?=$	Question Mark Over Equals Sign	$\stackrel{?}{=}$
$\bar{=}$	Vertical Bar Over Equals Sign	$\underline{=}$

➤ $\sqrt{5} \approx 2.236$

$\cdot=$ $\underset{\cdot}{=}$ $\overset{\cdot\cdot}{=}\underset{\cdot\cdot}{=}$ $\circ=$ $\triangle=$ $?=$ $\bar{=}$






➤ $5(3) + 2(-2) \cong 11$

$\cdot=$ $\underset{\cdot}{=}$ $\overset{\cdot\cdot}{=}\underset{\cdot\cdot}{=}$ $\circ=$ $\triangle=$ $?=$ $\bar{=}$

If a symbol is encountered which does not appear on this list, construct a symbol in accordance with the same principles. A transcriber-created symbol should be listed on the Special Symbols page.

➤ $A \triangleq B$ $\triangle=$ $\cdot=$ $\underset{\cdot}{=}$ $\overset{\cdot\cdot}{=}\underset{\cdot\cdot}{=}$ $\circ=$ $\triangle=$ $?=$ $\bar{=}$

The remainder of the list provided in the code contains other modified comparison signs most commonly used.

Modified Tilde		
	Dot Under Tilde	$\underset{\cdot}{\sim}$
	Dot Over Tilde	$\overset{\cdot}{\sim}$
Modified Horizontal Bar		
	Caret Over Bar	$\overset{\wedge}{-}$
	Caret Under Bar	$\underset{\wedge}{-}$
	("is perspective to")	
	Dot under Bar	$\underset{\cdot}{-}$

If the horizontal bar is modified by a dot *over* it, the combination is a modified sign of operation ("minus with dot over" signifying "proper difference"). The five-step rule is not used for this symbol. See Section 5.4.7.

Instructions: Use the five-step rule to show the horizontal grouping sign in the last item.

PRACTICE 12G

Modified Signs of Comparison and More

- $A \doteq 3.14r^2$
- $\angle b \doteq \frac{1}{2} \widehat{EB}$
- The symbol $\hat{=}$ is used to make a definition.
- $x \sim \mathcal{N}(0, 1)$
- $x^n = \underbrace{x \cdot x \cdot x \cdot \dots \cdot x}$

$$\gg \sum_{\substack{i,j,k \\ i < j < k}}$$



Analysis: The Sigma is modified directly under with letters i, j, k. That, in turn, is modified directly under with the expression i < j < k. Review the rules regarding use of the English-letter indicator in Lesson 3. An English-letter indicator is needed for the first letter "j" because it is a "single letter" according to the definition in the Nemeth Code. The other letters are either touching indicators or are preceded or followed by a sign of comparison, so an English-letter indicator is not needed.

12.13.1 Parallel Horizontal Bars. When two or more parallel horizontal bars are the same length and apply to exactly the same expression, they are treated as a single modifier. In such cases, the directly-over or the directly-under indicator is used only once, and the symbol for the bar is used as many times as necessary to correspond to the print text.

$$\gg \overline{9} \quad \dots$$

$$\gg (\overline{A}) \quad \dots$$

$$\gg \underline{x} \quad \dots$$

Attention: Two parallel horizontal bars must not be misinterpreted as the equals sign, and three parallel horizontal bars must not be misinterpreted as the identity sign. The symbol's meaning can be determined by reading the surrounding text for context. See Section 5.8 ("Signs of Comparison Compounded Vertically") to review equals sign printed above or below a sign of comparison.

12.14 Individual Modifiers

When two or more modifiers do not apply to exactly the same symbols but cover different portions of the same expression, the longer modification encloses the entire modified expression. Within the long expression, each inner expression is modified individually.

$$\gg \overline{\overline{\underline{x} \times \underline{x}}} \quad \dots$$

Analysis: In the simulated braille below, the longer modification indicators and symbols are highlighted; the inner modified expressions are underlined.



The five-step rule and the contracted form of transcription may be used together.

$$\Rightarrow \overline{(\bar{a}A + \bar{c}C)} \quad \cdot$$

Analysis: In the simulated braille below, the longer modification indicators and symbols are highlighted; the inner modified expressions are underlined.

$$\cdot \cdot$$

$$\Rightarrow \overline{P(\bar{x}^2)} \quad \cdot$$

Analysis: In the simulated braille below, the longer modification indicators and symbols are highlighted; the inner modified expression is underlined.

$$\cdot \cdot$$

Reminder: The subscript level indicator must be repeated before the modified expression indicator so the dot 5 is not misread as a baseline indicator.

12.15 Simultaneous Modifiers

When an expression is simultaneously modified both above and below, the modifier *below* is transcribed first and the modifier *above* is transcribed second. The termination indicator is used only at the end of the entire modification.

$$\Rightarrow \sum_3^7 \quad \cdot$$

$$\Rightarrow \prod_{k=2}^6 a_k \quad \cdot$$

Recall from 12.8 that, when the modifier is wider than the modified symbol (in this case, $k = 2$ is wider than Π), the print copy might insert extra space to clarify what exactly is being modified. In braille, the modified Pi is unspaced from the letter "a".

$$\Rightarrow \bar{x} \quad \cdot$$

$$\Rightarrow \overline{x + y} \quad \cdot$$


As you study the eight constructions in the next example, note the use or nonuse of the English-letter indicator and the use or nonuse of the numeric indicator. The simulated braille is in Nemeth. Code switch indicators are omitted.

Example 12-19

(1) $\sum_{k=1}^n (2k) = n(n + 1)$




(2) $\sum_{i,j=1}^n a_{ij}x_i x_j$




(3) $\sum_{x_1=0}^1 \sum_{x_2=0}^1$



(4) $\bigcup_{r=1}^n A_r = A_1 \cup A_2 \dots$



(5) $\prod_{x \in a} c^x$




(6) $\frac{b=2}{x+y} \frac{a=3}{}$



(7) $\sum_{i=1}^6 \overline{P_{i-1} P_i}$



$$(8) \quad (x + y)^n = \sum_{k=0}^n \binom{n}{k} x^{n-k} y^k$$



This is a linked expression where $(x + y)^n$ is the anchor and

$= \sum_{k=0}^n \binom{n}{k} x^{n-k} y^k$ is the link. Recall from Lesson 8 that when a linked

expression will not fit on one braille line, start a runover line with the link.

The equals sign within the modification is not a suitable division site because the components of an expression modified according to the five-step rule should not be divided between lines. (See [12.2](#).)

PRACTICE 12H

Expressions with More Than One Modifier

$$1. \quad \overline{\overline{9}} \cdot \overline{\overline{3}} = \overline{\overline{9 \cdot 3}}$$

$$5. \quad \overline{\overline{x_a + y^n}}$$

$$2. \quad \overline{\overline{A^n}}$$

$$6. \quad \overline{\overline{N}}$$

$$3. \quad \overline{\overline{A \cap B \cap C}}$$

$$7. \quad \sum_{i=1}^k \sum_{j=1}^k$$

$$4. \quad \overline{\overline{a_n + b_p}}$$

$$8. \quad \prod_{\substack{j=1 \\ j \neq k}}^n$$

Format

12.16 Formal Proof [NC Rule 26.7]

A proof is a valid argument that establishes the truth of a mathematical statement. It is often introduced by a heading such as *Theorem*, *Proposition*, or *Lemma*. A progressive sequence of statements leads to the conclusion. In a formal proof, every step of the argument is shown and each step is supported by a definition or by a previously proven statement.

Lesson 11 explained how to transcribe a mathematical statement. Those guidelines are summarized in steps a-c, below, in the context of a proof.

- a. A blank line precedes the beginning of the proof.
- b. *Heading*: The heading can be formatted as a paragraph heading or as a cell-5 or cell-7 heading, at the transcriber's discretion. Review Section 11.38.b for details.
- c. *Statement*: Continue with the text, using normal (3-1) paragraph style. When the statement is printed in a variant typeform and the proof follows, in regular type, it is recommended that the typeform be preserved for the statement in order to retain distinction.
- d. *Auxiliary Captions*: Paragraph headings such as Given, Hypothesis, Prove, or Conclusion begin in cell 3, without a blank line before the paragraph. Associated material follows the caption. Runovers go in cell 1.
- e. *Two-Column Proof*: See [12.16.1](#), below.
- f. *End of Proof Icon*: See [12.16.2](#), below.
- g. When the proof is complete, insert a blank line before continuing with the text.

12.16.1 Two-Column Proof. When a formal proof is presented by numbered steps printed in two columns, the layout is changed as follows.

- a. If there is a caption such as "Proof", follow the same pattern established in [step d](#), above ("Auxiliary Captions").
- b. The column format is changed to a list in braille. A transcriber's note must call attention to the change in format. See [step c](#), below, for a sample transcriber's note.

A blank line is inserted before the list. If there are column headings, such as "Statement" and "Reason", see [step c](#), below. Each step begins in cell 1, starting with the first item from the left column. Runovers are in cell 3. The related item from the right column begins in cell 1 on the next line, with runovers in cell 3.

Each item must be labeled with an identifier as described below.

- c. *Step Identifiers*: Typically, the print copy includes the column headings "Statements" and "Reasons". In braille, the column headings are replaced with a letter – "S" or "R", respectively – as part of each step number. For example, "1S" is Statement 1, followed on the next line by "1R" which is Reason 1. If other column headings appear in print, the transcriber chooses an appropriate letter to represent each heading. If no column headings appear in print, use "S" for the left-column items and "R" for the right-column items.

If the steps are not numbered in print, identifiers are added in the manner stated above. A transcriber's note must specify the meaning of the letter labels. Any transcriber-added numbers or letters must also be explained.

Sample transcriber's note: "Proofs printed with steps in columns headed *Statements* and *Reasons* are transcribed with an S or R immediately following the step number to show the column in which the step appears. Each step from the *Statements* column is immediately followed by the corresponding step from the *Reasons* column."

Identifiers are transcribed in either UEB or Nemeth according to code-switching guidelines.


- 12.16.2 **End of Proof Icon**. Sometimes a shape (often a bold rectangle or square) or the Latin abbreviation "Q.E.D." is used to denote the end of a proof. The shape is often printed at the right margin and may go unnoticed. Watch for it. Regardless of the shape or abbreviation used in print, the icon shown below is used in braille, and is transcribed following the last word or character in the proof, preceded by one blank cell. The icon is created using the UEB transcriber-defined shape indicator and may be used in either Nemeth or UEB without switching. The icon must be listed on the Special Symbols page. (Details regarding a Special Symbols page in a UEB with Nemeth transcription will be covered in the Final Lesson.)



If the shape indicates the end of a long example or exercise, the symbol for the filled-in or shaded shape is used. See Section 11.6 in Lesson 11.

Example 12-20

The end of proof mark **■** is used in mathematics to denote the end of a proof, in place of the traditional abbreviation "Q.E.D." for the Latin phrase "**quod erat demonstrandum**", meaning "which was to be shown".



Example 12-21

(Assume the required transcriber's note regarding the two-column proof appears on the Transcriber's Notes page.)

THEOREM 2. All right angles are equal.

Given: $\angle ABC$ and $\angle DEF$ are right angles.

Prove: $\angle ABC$ equals $\angle DEF$.

<i>Statements</i>	<i>Reasons</i>
1. $\angle ABC$ and $\angle DEF$ are right angles.	1. Given.
2. $\angle ABC = 90^\circ$, $\angle DEF = 90^\circ$.	2. A right angle contains 90 degrees.
3. $\angle ABC = \angle DEF$.	3. Transitivity postulate. ■

7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23

Line 7: A blank line precedes the proof. Box lines are not required.

Line 8: The heading is printed in capital letters and also in a nonregular typeface (boldface).

When a paragraph heading is printed in capital letters, typeform is disregarded.

Lines 8-9: The paragraph style is 3-1. Distinctive typeform (boldface) is retained in the statement.

Lines 10-11 and 12-13: Each auxiliary caption follows print regarding typeform (italics, in this example), and uses a 3-1 paragraph style.

Line 14: A blank line precedes the list.

Lines 15-22: Each item in the 2-column proof begins in cell 1, with runovers in cell 3. Identifying letters S and R are combined with each step number.

Line 22: A dark square is printed in the right margin to mark the end of the proof. The "qed" icon is transcribed.

Line 23: A blank line follows the proof.

PRACTICE 12I

Instructions: Create a Transcriber's Notes page that might appear in a volume which contains the proof shown in Practice 12J. A statement citing the code book is required in every volume that uses Nemeth. In the first paragraph, state the title and edition of the Nemeth Code as well as any BANA updates. Something like this:

Mathematical content is transcribed according to *The Nemeth Braille Code for Mathematics and Science Notation, 2022*.

In the second paragraph, explain the step-number format as described in 12.16.1.c. Refer to *Braille Formats* for further guidelines regarding the structure of a Transcriber's Notes page.

PRACTICE 12J

Given: $3x = 7 - \frac{1}{2}x$

To Prove: $x = 2$

STEP	REASON
1. $3x = 7 - \frac{1}{2}x$	1. GIVEN
2. $6x = 14 - x$	2. Multiplication Property
3. $7x = 14$	3. Addition Property
4. $x = 2$	4. Division Property

For further practice, see Appendix A—Reading Practice.

EXERCISE 12

Prepare Exercise 12 for your grader.

PRACTICE 12H

1
2
3
4
5
6
7
8
9
10
11

The page contains 11 rows of Braille characters for reading practice. Row 1 contains five groups of six Braille cells. Row 2 is a single continuous line of 100 dotted Braille cells. Row 3 contains two groups of two Braille cells. Row 4 contains four groups of six Braille cells. Row 5 contains two groups of six Braille cells. Row 6 contains two groups of six Braille cells. Row 7 contains two groups of six Braille cells. Row 8 contains two groups of six Braille cells. Row 9 contains two groups of six Braille cells. Row 10 contains seven groups of six Braille cells. Row 11 contains seven groups of six Braille cells.

