

LESSON 17

- [SYSTEM OF EQUATIONS](#)
 - [Enlarged Signs of Grouping](#)
- [DETERMINANTS AND MATRICES](#)

[Answers to Practice Material](#)

LESSON PREVIEW

Enlarged grouping signs are often encountered in the topics studied in this lesson: systems of equations and arrays (matrices and determinants). Format rules are given for these spatial arrangements, including further considerations regarding commentary.

SYSTEM OF EQUATIONS

[NC 25.9]

17.1 Definition and Recognition

A system of equations, sometimes called "simultaneous equations," is a collection of two or more equations which share variables. The student identifies the value of each variable by "solving the system." You can recognize a system by noticing the arrangement of equations on two or more lines. If the system consists of two equations, there will be two variables to solve (typically x and y); if the system consists of three equations, there will usually be three variables to solve (typically x , y , and z). The equations may or may not be joined by an enlarged grouping sign.

Here is a sample of a system of two equations using variables x and y .

$$\begin{array}{l} 4x - y = 10 \\ 2x = 12 - 3y \end{array}$$

17.2 Transcription Rules for Systems of Equations

A system of equations is a spatial arrangement and is transcribed as follows.

- (a) One blank line is left above and below the system. Placement of code switch indicators follows the general rules for spatial arrangements.
- (b) Use and nonuse of the numeric indicator follows Nemeth rules for nonspatial material, even though this is a spatial arrangement. (See also [subitem \(d\)](#), below.)
- (c) Alignment is maintained only if terms and symbols are aligned in print. If terms are aligned on one side of the equals sign but not on the other side, follow print. Alignment is disregarded if only the equals signs are aligned.
- (d) When a space is inserted within the equation for the purpose of maintaining alignment, a numeric indicator is not used.

The same rules apply to a system of inequalities. (See [Example 17-4](#)).

Note that, if a separation line is present, it is a spatially arranged addition problem and Nemeth rules for spatial addition are followed. (See Lesson 9.)

PRACTICE 17A

1. Solve and check:

$$2x + 3y = 2$$

$$8x - 4z = 3$$

$$3y - 8z = -1$$

2. Solve:

$$2x - 5y + 6z = 11$$

$$3x - 2y + 3z = 9$$

$$2x + 4y - 9z = -3$$

3. Add:

$$3x - y = 7$$

$$\underline{2x + y = 8}$$

$$5x + 0 = 15$$

Enlarged Signs of Grouping

[NC 19.6]

17.3 A Unified Expression

The Nemeth Code calls an arrangement a "unified expression" when equations are grouped together using an enlarged grouping symbol. Here is a print sample of a unified system of equations, using a left enlarged brace.

$$\left\{ \begin{array}{l} x = y \\ 5x - y = 4 \end{array} \right.$$

17.4 Transcription Rules for Enlarged Signs of Grouping

- If only the left or right grouping sign is shown in print, only that symbol is shown in braille.
- Enlarged grouping symbols are transcribed on each line of the unified expression and are vertically aligned.
- When terms are not aligned in print, each line of the unified expression begins in the same cell. When there is a left enlarged grouping symbol, each expression will begin in the cell which immediately follows the left enlarged grouping symbol.

- (d) When the expressions require vertical alignment, at least one item must begin in the cell which immediately follows the left enlarged grouping symbol. The numeric indicator is not required before the first numeric character following a left grouping symbol.
- (e) At least one item ends in the cell which immediately precedes the right enlarged grouping symbol.

17.4.1 **Left Enlarged Brace.** The left enlarged brace curves and points to the left in print. Notice that the enlarged braille symbol is formed by inserting a dot 6 before the second cell of the normal brace symbol.

⠠⠨	Left Brace (normal size)	{
⠠⠠⠨	Left Enlarged Brace (covering two or more lines)	

Example 17-5

$$\begin{cases} x = y \\ 5x - y = 4 \end{cases}$$

Print observation: A left enlarged brace groups the unified system of equations. Terms are not vertically aligned.

1	⠠⠨	
2	⠠⠠⠨	
3	⠠⠠⠨	⠠⠠⠨
4	⠠⠠⠨	⠠⠠⠨
5	⠠⠠⠨	⠠⠠⠨
6	⠠⠨	

Braille observation: The enlarged grouping symbols are aligned. The terms are not aligned in print; each equation begins in the cell following the left enlarged brace. A numeric indicator is not needed for the numeral 5 because it is not preceded by a space. A numeric indicator is required for the numeral 4.

17.4.2 **Right Enlarged Brace.** The right enlarged brace curves and points to the right in print. Notice that the enlarged braille symbol is formed by inserting a dot 6 before the second cell of the normal brace symbol.

⠠⠨	Right Brace (normal size)	}
⠠⠠⠨	Right Enlarged Brace (covering two or more lines)	

Lines 1-2: The paragraph begins in cell 3. The runover is in cell 1.
 Line 3: A blank precedes the spatial arrangement that is embedded in the paragraph.
 Line 4: The narrative continues, in the runover cell of the paragraph. The opening switch is placed on this line only. The narrative continues after the top right enlarged grouping symbol.
 Line 5: The second line of the embedded spatial arrangement is aligned with the line above, starting in the same cell and with the right grouping symbols aligned.
 Line 6: A blank follows the spatial arrangement that lies within the paragraph.
 Line 7: The paragraph continues.

PRACTICE 17B

After solving for x and y , we determine that the system $\begin{cases} 2x - 3y = 17 \\ 3x + 2y = 6 \end{cases}$ has the solution set $\{(4, -3)\}$. Name the solution set for this system.

$$\begin{cases} x + 2y = 6 \\ 2x - y = 7 \end{cases}$$

17.6 Enlarged Parentheses

Notice that the enlarged braille parentheses are formed by inserting a dot 6 before the normal parenthesis symbol.

⠠	Left Parenthesis (normal)	(
⠠⠠	Left Enlarged Parenthesis (covering two or more lines)	
⠡	Right Parenthesis (normal))
⠡⠡	Right Enlarged Parenthesis (covering two or more lines)	


17.7 Placement of Symbols

Symbols which appear outside of the enlarged grouping symbol and which apply to the arrangement are placed on the top line of the arrangement, even if the items are centered in print.

Example 17-8

$$x = \left(\begin{array}{l} j = lkjkl \\ = l \\ kjkljlkjkl = k \end{array} \right.$$

Print observation: A three-line system is grouped on the left with an enlarged parenthesis. The anchor, x, is centered with respect to the enlarged grouping symbol to its right. The equals signs in the grouped arrangement are aligned, but the terms are not.



Braille observation: The anchor is aligned with the top line of the arrangement. Each enlarged grouping symbol begins in the same cell. The grouped items are left adjusted because the terms are not aligned. A general omission symbol denotes the blank space before the middle equals sign. English-letter indicators are not needed for the single letters x, j, l, and k because they are next to a sign of comparison. Code switch indicators do not interfere with the spatial arrangement.


17.8 Placement of Identifiers and Punctuation

Identifiers and punctuation which appear outside of the enlarged grouping symbol and which apply to the arrangement are placed on the top line of the arrangement, even if the items are centered in print. [Example 17-9](#) uses the same expression from [Example 17-8](#), now preceded by an identifier. An itemized spatial arrangement is not considered to be embedded. The code switch indicators are placed outside of the arrangement.

Example 17-9

$$15. x = \left(\begin{array}{l} j = lkjkl \\ = l \\ kjkljlkjkl = k \end{array} \right.$$

Print observation: The item number and the anchor are centered with respect to the associated enlarged grouping symbol to the right.




Braille observation: Item number and anchor are aligned with the top line of the arrangement. The code switch indicators are placed outside of the arrangement.

Example 17-10

... such that
$$\left. \begin{array}{l} \frac{1}{2}x + y = 7 \\ 5x - y = 4 \end{array} \right\} \text{. First, solve for } y \text{ in terms of } x.$$

Print observation: A vertical arrangement is embedded within the narrative. It is grouped on the right with a right enlarged parenthesis. A period follows the enlarged parenthesis, centered to the arrangement. The narrative continues on the same line.



Braille observation:

Line 1: A blank line precedes the line with the embedded spatial arrangement.

Line 2: The top line of the arrangement begins on the main line of the text. The code switch indicators are also placed on this line. The paragraph continues after the period.

Line 3: The second line of the arrangement is placed here. The right enlarged parenthesis is aligned with the identical symbol on the line above. No code switch indicators are placed in this line.

Line 4: A blank line follows the embedded spatial arrangement.

Line 5: The paragraph text continues.

Braille observation: The left enlarged brace is transcribed as printed, beginning in cell 3 (displayed to narrative). The transcriber-inserted grouping symbol is placed between the arrangement and the conditions. The Nemeth Code terminator is placed in cell 1 after the blank line.

17.10.5 Line-by-Line Commentary. When comments refer to individual lines in a grouped expression, the method presented in Lesson 16 is applied, indenting the comments four cells to the right of the of the runover position of the line to which they apply. Because Nemeth enlarged grouping symbols are used, the comments are uncontracted. A transcriber's note is required to explain the layout.

Example 17-16

This system of equations models two bikers' routes, where x represents time (in hours) and y represents distance (in miles).

$$\left\{ \begin{array}{ll} y = 1.5x & \text{Su's route} \\ y = 12 - 2.5x & \text{Jayla's route} \end{array} \right.$$

1 $y = 1.5x$ Su's route

2 $y = 12 - 2.5x$ Jayla's route

3

4

5

6

7

8

9

10

11

12

13

14

Lines 4-7: The transcriber's note reads as follows. "Comments printed beside related math problems are placed on the line following the expression, blocked four cells to the right of the runover position of the expression."

Lines 8 and 13: A blank line precedes and follows the spatially-arranged material.

PRACTICE 17D

Find x and y in terms of a and b .

1.
$$\begin{cases} ax + by = 0 \\ x + y = 1 \end{cases} \quad (a \neq b)$$

2.
$$\begin{cases} ax + by = 0 \\ a^2x + b^2y = 1 \end{cases} \quad (a \neq 0, b \neq 0, a \neq b)$$

3.
$$\left. \begin{array}{l} a = \frac{x+y}{x-y} \\ b = \frac{x-y}{x+y} \end{array} \right\} -1 < x < 1, -1 < y < 1$$

4. Solve.

$$\left. \begin{array}{l} 11z - 4z + 2z = 27 \\ 7z + 2z = 27 \\ 9z = 27 \\ z = 3 \end{array} \right\} \text{Combine like terms.}$$


17.11 More Enlarged Signs of Grouping [NC Rule 19]

In addition to the enlarged braces and parentheses, the Nemeth Code provides symbols for six other enlarged signs of grouping. Notice that each enlarged braille symbol is formed by inserting a dot 6 before the ⠠, ⠡, or ⠢ symbol of the normal-sized grouping symbol.

Vertical Bar		
⠠⠼	Single, Normal	
⠠⠠⠼	Double, Normal	
⠠⠠⠠⠼	Single, Enlarged	
⠠⠠⠠⠠⠼	Double, Enlarged	
Barred Brace		
⠠⠠⠠⠼	Left, Normal	{
⠠⠠⠠⠼	Right, Normal	}
⠠⠠⠠⠠⠼	Left, Enlarged	
⠠⠠⠠⠠⠼	Right, Enlarged	
Bracket		
⠠⠠⠼	Left, Normal	[
⠠⠠⠼	Right, Normal]
⠠⠠⠠⠼	Left, Enlarged	
⠠⠠⠠⠼	Right, Enlarged	
Angle Bracket		
⠠⠠⠠⠼	Left, Normal	<
⠠⠠⠠⠼	Right, Normal	>
⠠⠠⠠⠠⠼	Left, Enlarged	
⠠⠠⠠⠠⠼	Right, Enlarged	


Barred Bracket		
	Left, Normal	[
	Right, Normal]
	Left, Enlarged	
	Right, Enlarged	
Half Bracket		
	Upper Left, Normal	[
	Upper Right, Normal]
	Upper Left, Enlarged	
	Upper Right, Enlarged	
	Lower Left, Normal	L
	Lower Right, Normal	J
	Lower Left, Enlarged	
	Lower Right, Enlarged	

Example 17-17



Instructions: Follow "top alignment" rules for these side-by-side arrays. Place the leftmost character in cell 1.

PRACTICE 17E



DETERMINANTS AND MATRICES

[NC 25.8]

17.12 Definition and Recognition

Determinants and matrices (singular: matrix) are arrangements of items in rows and columns which are enclosed between left and right grouping symbols. Items can be numbers, symbols, or mathematical expressions.

Here is a 2×2 ("two by two") determinant enclosed between enlarged vertical bars.

$$\left| \begin{array}{cc} a & b \\ c & d \end{array} \right|$$

Here is a 2×3 ("two by three") matrix enclosed between enlarged brackets.

$$\left[\begin{array}{ccc} 1 & 9 & -13 \\ 20 & 5 & -6 \end{array} \right]$$

A matrix can also be composed of only one column or only one row. Row matrices will be studied at the end of this section.

Determinants and matrices may also be referred to as "arrays."

17.13 Transcription Rules for Determinants and Matrices

Observe the following rules regarding the transcription of the two samples shown in [Section 17.12](#).

- 17.13.1 **Blank Lines.** Determinants and matrices are spatial arrangements. Thus, a blank line is to be left above and below each array. If the arrangement begins at the top of a braille page, it may begin on line 1 provided no running head is in use; if the arrangement ends at the bottom of a braille page, it may end on line 25. In either case, the rightmost symbol of the arrangement must not lie within three cells of the page number. (See Lesson 9 for details regarding layout of a spatial arrangement at the top or bottom of a page.)
- 17.13.2 **Grouping Symbols.** For an array consisting of two or more rows, enlarged grouping symbols are used on each braille line. Grouping symbols are vertically aligned.
- 17.13.3 **Alignment and Spacing of Items.** Each entry is moved as far left as possible in its column. Consequently, each left grouping symbol will be in direct contact with the first entry of each row in the array.

One column of blank cells is left between the columns of the arrangement. That is, one blank cell separates the widest entry in a column from the beginning of the next column. Even if entries contain a space, only one space is left between columns. See [Example 17-20](#).

PRACTICE 17F

1. $\begin{vmatrix} 1 & 2 \\ 2 & -1 \end{vmatrix}$

2. $\begin{pmatrix} 1 & -\frac{3}{4} & \frac{5}{3} \\ 2 & 5 & 12 \end{pmatrix}$

3. $\begin{bmatrix} a & b & c \\ 0 & 0 & 0 \end{bmatrix}$

4. $\begin{vmatrix} ab & cd \\ ac & ce \end{vmatrix}$

5. Explain why points (a_1, b_1) , (a_2, b_2) , and (a_3, b_3) are collinear if and only if

$$\begin{vmatrix} a_1 & b_1 & 1 \\ a_2 & b_2 & 1 \\ a_3 & b_3 & 1 \end{vmatrix} = 0$$

PRACTICE 17G

- A. The unit vectors of a three dimensional Cartesian coordinate system are shown below.

$$\hat{\mathbf{i}} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \hat{\mathbf{j}} = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \text{ and } \hat{\mathbf{k}} = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}.$$

17.15 Ellipses, Single Dots, and Blank Entries

The following symbols represent ellipses, a single dot, or a blank entry in a matrix or a determinant.

$\cdot\cdot$	Blank Entry	
$\cdot\cdot\cdot$	Ellipsis, Diagonal,	$\cdot\cdot$
	lower left to upper right	
$\cdot\cdot\cdot$	Ellipsis, Diagonal,	$\cdot\cdot$
	upper left to lower right	
$\cdot\cdot\cdot$	Ellipsis, Horizontal	\dots
$\cdot\cdot\cdot$	Ellipsis, Vertical	\vdots
\cdot	Single Dot	\cdot

Each symbol is positioned as far left as possible in its column. A transcriber’s note must explain the use of the short dash to represent the blank space. Sample transcriber’s note:

A short dash represents a blank entry.

Example 17-24

$$\begin{vmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdot & \cdot & \cdots & \cdot \\ a_{n1} & a_{n2} & \cdots & a_{nn} \end{vmatrix}$$

Print observation: A horizontal ellipsis is in each entry in column 3. A single dot is printed in row 3, columns 1, 2, and 4.

1	$\cdot\cdot$	
2	$\cdot\cdot\cdot$	
3	$\cdot\cdot\cdot$	$\cdot\cdot$
4	$\cdot\cdot\cdot$	$\cdot\cdot$
5	\cdot	\cdot
6	$\cdot\cdot\cdot$	$\cdot\cdot$
7	$\cdot\cdot\cdot$	\dots
8	$\cdot\cdot$	\cdot

Braille observation: A baseline indicator is needed to assure the right grouping symbols on rows 1, 2, and 4 are on the same level as the left grouping symbols.

PRACTICE 17H

In the next equation, matrix Y is expressed as the product of matrices B and X.

$$\begin{bmatrix} y_1 \\ y_2 \\ y_3 \\ \vdots \\ y_r \end{bmatrix} = \begin{bmatrix} b_{11} & b_{12} & \cdots & b_{1n} \\ b_{21} & b_{22} & \cdots & b_{2n} \\ b_{31} & b_{32} & \cdots & b_{3n} \\ \vdots & \vdots & \ddots & \vdots \\ b_{r1} & b_{r2} & \cdots & b_{rn} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ \vdots \\ x_n \end{bmatrix}$$

Instructions: Place a transcriber's note before item 2 explaining the treatment of the blank entries.

PRACTICE 17I

1. $A = \begin{bmatrix} a_{11} & \cdots & a_{1M} \\ \vdots & \ddots & \vdots \\ a_{K1} & \cdots & a_{KM} \end{bmatrix}$

2. $B = \begin{bmatrix} b_{11} & b_{12} & \cdots \\ \vdots & \ddots & \\ b_{K1} & & b_{KK} \end{bmatrix}$

1	$\frac{1}{2}$	$\frac{1}{3}$	$\frac{1}{4}$	$\frac{1}{5}$	$\frac{1}{6}$	$\frac{1}{7}$	$\frac{1}{8}$	$\frac{1}{9}$	$\frac{1}{10}$	$\frac{1}{11}$
2		$\frac{1}{2}$	$\frac{1}{3}$	$\frac{1}{4}$	$\frac{1}{5}$	$\frac{1}{6}$	$\frac{1}{7}$	$\frac{1}{8}$	$\frac{1}{9}$	$\frac{1}{10}$
3	$\frac{1}{2}$	$\frac{1}{3}$	$\frac{1}{4}$	$\frac{1}{5}$	$\frac{1}{6}$	$\frac{1}{7}$	$\frac{1}{8}$	$\frac{1}{9}$	$\frac{1}{10}$	$\frac{1}{11}$
4		$\frac{1}{2}$	$\frac{1}{3}$	$\frac{1}{4}$	$\frac{1}{5}$	$\frac{1}{6}$	$\frac{1}{7}$	$\frac{1}{8}$	$\frac{1}{9}$	$\frac{1}{10}$
5			$\frac{1}{2}$	$\frac{1}{3}$	$\frac{1}{4}$	$\frac{1}{5}$	$\frac{1}{6}$	$\frac{1}{7}$	$\frac{1}{8}$	$\frac{1}{9}$
6				$\frac{1}{2}$	$\frac{1}{3}$	$\frac{1}{4}$	$\frac{1}{5}$	$\frac{1}{6}$	$\frac{1}{7}$	$\frac{1}{8}$
7					$\frac{1}{2}$	$\frac{1}{3}$	$\frac{1}{4}$	$\frac{1}{5}$	$\frac{1}{6}$	$\frac{1}{7}$
8						$\frac{1}{2}$	$\frac{1}{3}$	$\frac{1}{4}$	$\frac{1}{5}$	$\frac{1}{6}$
9							$\frac{1}{2}$	$\frac{1}{3}$	$\frac{1}{4}$	$\frac{1}{5}$
10	$\frac{1}{2}$	$\frac{1}{3}$	$\frac{1}{4}$	$\frac{1}{5}$	$\frac{1}{6}$	$\frac{1}{7}$	$\frac{1}{8}$	$\frac{1}{9}$	$\frac{1}{10}$	$\frac{1}{11}$
11										

One blank cell separates the widest entry in a column, including any runovers, from the beginning of the next column.

17.17.2 **Runovers Without Indentation.** If the technique shown in [Section 17.17.1](#) is not feasible, entries may be run over to new lines without indentation. Preference rules for runovers of mathematical expressions need not be observed if space would be saved. In order to distinguish each row, a blank line is inserted between them. Enlarged grouping symbols are transcribed on those blank lines within the arrangement.

Example 17-30

1) ... shown below.

$$\mathbf{AB} = \begin{bmatrix} a_{11}b_{11} + a_{12}b_{21} & a_{11}b_{12} + a_{12}b_{22} & a_{11}b_{13} + a_{12}b_{23} \\ a_{21}b_{11} + a_{22}b_{21} & a_{21}b_{12} + a_{22}b_{22} & a_{21}b_{13} + a_{22}b_{23} \\ a_{31}b_{11} + a_{32}b_{21} & a_{31}b_{12} + a_{32}b_{22} & a_{31}b_{13} + a_{32}b_{23} \end{bmatrix}$$

$$\begin{array}{r}
 1 \quad \left[\begin{array}{cccc} e^{2t} & e^{2t} & e^{2t} & e^{2t} \end{array} \right] \\
 2 \quad \left[\begin{array}{cccc} e^{2t} & e^{2t} & e^{2t} & e^{2t} \end{array} \right] \\
 3 \quad \left[\begin{array}{cccc} e^{2t} & e^{2t} & e^{2t} & e^{2t} \end{array} \right] \\
 4 \quad \left[\begin{array}{cccc} e^{2t} & e^{2t} & e^{2t} & e^{2t} \end{array} \right] \\
 5 \quad \left[\begin{array}{cccc} e^{2t} & e^{2t} & e^{2t} & e^{2t} \end{array} \right] \\
 6 \quad \left[\begin{array}{cccc} e^{2t} & e^{2t} & e^{2t} & e^{2t} \end{array} \right] \\
 7 \quad \left[\begin{array}{cccc} e^{2t} & e^{2t} & e^{2t} & e^{2t} \end{array} \right] \\
 8 \quad \left[\begin{array}{cccc} e^{2t} & e^{2t} & e^{2t} & e^{2t} \end{array} \right] \\
 9 \quad \left[\begin{array}{cccc} e^{2t} & e^{2t} & e^{2t} & e^{2t} \end{array} \right] \\
 10 \quad \left[\begin{array}{cccc} e^{2t} & e^{2t} & e^{2t} & e^{2t} \end{array} \right] \\
 11 \quad \left[\begin{array}{cccc} e^{2t} & e^{2t} & e^{2t} & e^{2t} \end{array} \right] \\
 12 \quad \left[\begin{array}{cccc} e^{2t} & e^{2t} & e^{2t} & e^{2t} \end{array} \right] \\
 13 \quad \left[\begin{array}{cccc} e^{2t} & e^{2t} & e^{2t} & e^{2t} \end{array} \right]
 \end{array}$$

17.17.3 **Runovers and Level Indicators.** As noted in examples throughout this section, a baseline indicator is needed when a row entry in the rightmost column ends at a level other than the baseline, and it touches the right grouping symbol (subscripts which don't require a subscript indicator excepted). When a row entry requires a runover, this rule applies only to the last symbol of the entry. A baseline indicator is not used before a right enlarged grouping symbol if the expression which touches the right grouping symbol is continued on the next line.

Example 17-31

$$\left[\begin{array}{ccc} -2te^t + e^{2t} & (3t + 2)e^t - 2e^{2t} & -(2t + 1)e^t + 2e^{2t} \\ -2(t + 1)e^t + 2e^{2t} & (3t + 5)e^t - 4e^{2t} & -(2t + 2)e^t + 2e^{2t} \end{array} \right]$$

$$\begin{array}{r}
 1 \quad \left[\begin{array}{ccc} -2te^t + e^{2t} & (3t + 2)e^t - 2e^{2t} & -(2t + 1)e^t + 2e^{2t} \\ -2(t + 1)e^t + 2e^{2t} & (3t + 5)e^t - 4e^{2t} & -(2t + 2)e^t + 2e^{2t} \end{array} \right] \\
 2 \quad \left[\begin{array}{ccc} -2te^t + e^{2t} & (3t + 2)e^t - 2e^{2t} & -(2t + 1)e^t + 2e^{2t} \\ -2(t + 1)e^t + 2e^{2t} & (3t + 5)e^t - 4e^{2t} & -(2t + 2)e^t + 2e^{2t} \end{array} \right] \\
 3 \quad \left[\begin{array}{ccc} -2te^t + e^{2t} & (3t + 2)e^t - 2e^{2t} & -(2t + 1)e^t + 2e^{2t} \\ -2(t + 1)e^t + 2e^{2t} & (3t + 5)e^t - 4e^{2t} & -(2t + 2)e^t + 2e^{2t} \end{array} \right] \\
 4 \quad \left[\begin{array}{ccc} -2te^t + e^{2t} & (3t + 2)e^t - 2e^{2t} & -(2t + 1)e^t + 2e^{2t} \\ -2(t + 1)e^t + 2e^{2t} & (3t + 5)e^t - 4e^{2t} & -(2t + 2)e^t + 2e^{2t} \end{array} \right] \\
 5 \quad \left[\begin{array}{ccc} -2te^t + e^{2t} & (3t + 2)e^t - 2e^{2t} & -(2t + 1)e^t + 2e^{2t} \\ -2(t + 1)e^t + 2e^{2t} & (3t + 5)e^t - 4e^{2t} & -(2t + 2)e^t + 2e^{2t} \end{array} \right] \\
 6 \quad \left[\begin{array}{ccc} -2te^t + e^{2t} & (3t + 2)e^t - 2e^{2t} & -(2t + 1)e^t + 2e^{2t} \\ -2(t + 1)e^t + 2e^{2t} & (3t + 5)e^t - 4e^{2t} & -(2t + 2)e^t + 2e^{2t} \end{array} \right] \\
 7 \quad \left[\begin{array}{ccc} -2te^t + e^{2t} & (3t + 2)e^t - 2e^{2t} & -(2t + 1)e^t + 2e^{2t} \\ -2(t + 1)e^t + 2e^{2t} & (3t + 5)e^t - 4e^{2t} & -(2t + 2)e^t + 2e^{2t} \end{array} \right] \\
 8 \quad \left[\begin{array}{ccc} -2te^t + e^{2t} & (3t + 2)e^t - 2e^{2t} & -(2t + 1)e^t + 2e^{2t} \\ -2(t + 1)e^t + 2e^{2t} & (3t + 5)e^t - 4e^{2t} & -(2t + 2)e^t + 2e^{2t} \end{array} \right]
 \end{array}$$


Lines 3-6: Note that this example uses full margins in addition to applying runover techniques.
 Lines 3 and 5: A baseline indicator is not needed between the superscript "t" and the right grouping symbol because the expression continues on the next line.

Lines 4 and 6: The runover line of each row entry ends with the letter *t* in the superscript position. A baseline indicator is required before transcribing the right grouping symbol which follows, unspaced.

17.17.4 **Fractions in Arrays.** Fractions in an array may be transcribed linearly but may be arranged spatially if linear fractions take up too much space.

Example 17-32

$$\begin{bmatrix} \frac{1}{6} & \frac{5}{6} & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & \frac{1}{6} & 0 & \frac{5}{6} \\ \frac{5}{6} & 0 & 0 & \frac{1}{6} \end{bmatrix}$$




Observe the following details when transcribing the fractions spatially.

- A line is skipped between the rows containing the spatial fraction. Enlarged grouping symbols are transcribed on those blank lines within the arrangement.
- Operation signs and variables are placed on the same line as the spatial fraction line.
- Each entry is moved as far up as possible in its row. This includes entries that are not fractions.
- If the last item in a row is a superscript, or a subscript that uses a subscript indicator, a return to the baseline must occur before the right grouping symbol is transcribed. If the row extends fully to the grouping symbol, a baseline indicator is required to return to the baseline. If the row does not extend to the right grouping symbol, the space returns the reader to the baseline.

Example 17-33

$$\begin{vmatrix} \frac{3}{8}e^{2x} & -\frac{3}{8} & +\frac{1}{4} & -\frac{3}{4}x^2 \\ -\frac{3}{8}e^{2x} & +\frac{3}{8} & +\frac{3}{4} & +\frac{3}{4}x^2 \\ 3e^{2x} & -\frac{3}{8} & -\frac{3}{4} & +\frac{3}{4}x^2 \end{vmatrix}$$



The single-line entry in row 3, column 1, is transcribed on the top line of the row even though it is centered in its row in print.

17.17.5 **Keying.** When the array will not fit on the page even after trying the methods described in this section, a key may be devised. The technique of keying will be discussed in the next lesson.

Instructions: Use spatial fractions.

PRACTICE 17J

The derivative of T can be expressed as follows.

$$dT = \begin{bmatrix} \frac{\partial y_1}{\partial x_1} & \frac{\partial y_1}{\partial x_2} & \dots & \frac{\partial y_1}{\partial x_n} \\ \dots & \dots & \dots & \dots \\ \frac{\partial y_m}{\partial x_1} & \frac{\partial y_m}{\partial x_2} & \dots & \frac{\partial y_m}{\partial x_n} \end{bmatrix}$$

Instructions: Use runovers with indentation. Review spacing rules and "keep together" rules for abbreviated function names in Lesson 14.

PRACTICE 17K

In this Jacobian determinant,

$$\begin{vmatrix} 0 & 5 & 0 \\ 8x_1 & -2x_3 \cos(x_2x_3) & -2x_2 \cos(x_2x_3) \\ 0 & x_3 & x_2 \end{vmatrix} = -8x_1 \begin{vmatrix} 5 & 0 \\ x_3 & x_2 \end{vmatrix} = -40x_1x_2$$


the orientation of the resulting object is reversed.

17.18 Row Matrix

A row matrix has only one row. Because it is a matrix, it is transcribed as a spatial arrangement—that is, a line is left above and below the arrangement. Regular grouping symbols are used, not enlarged symbols, despite the larger appearance of the grouping symbols in print. Here is an example of a 1×5 row matrix.

Example 17-34

8. $[2\ 0\ 4\ 5\ 9]$




This is not an enclosed list because there are no commas between items. Assume that context will make it clear that this is a row matrix. A blank line precedes and follows the matrix. The brackets are tall in print, but are transcribed as regular brackets.

17.19 Embedded Arrays

If the array is embedded within narrative text, the required blank line is inserted on the line before and the line after the line where the arrangement lies. Text is placed on the same line as the array if it fits.

Example 17-35

Now we will look at a row matrix—a matrix consisting of just one row. $[2\ 0\ 5]$ is a 1×3 row matrix. Where have we seen a 1×5 row matrix?



A blank line precedes and follows the line which contains the embedded row matrix. The brackets are tall in print, but are transcribed as regular brackets.

PRACTICE 17L

If $A = \begin{bmatrix} 2 & -1 \\ 3 & 4 \end{bmatrix}$ and $B = \begin{bmatrix} 3 & -9 & 2 \\ 5 & 7 & -6 \end{bmatrix}$, find AB . The answer is

$$AB = \begin{bmatrix} 1 & -25 & 10 \\ 29 & 1 & -18 \end{bmatrix}.$$

PRACTICE 17M

1) Here are three examples of matrix operations.

a) $\begin{bmatrix} 5 & 4 \end{bmatrix} + \begin{bmatrix} 20 & 30 \end{bmatrix} = \begin{bmatrix} 25 & 34 \end{bmatrix}$

b) $\begin{bmatrix} 1 & 0 \\ 0 & 2 \\ 1 & 3 \end{bmatrix} + \begin{bmatrix} 2 & 4 \\ 1 & 5 \\ 0 & 6 \end{bmatrix} = \begin{bmatrix} 3 & 4 \\ 1 & 7 \\ 1 & 9 \end{bmatrix}$

c) $\begin{bmatrix} 2 & 1 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} = \begin{bmatrix} -1 & 2 \end{bmatrix}$

17.20 Use of Tactile Graphics for Enlarged Grouping Signs

Enlarged grouping symbols may be drawn in place of the braille equivalents, especially when space saving is a factor. Refer to BANA's *Guidelines and Standards for Tactile Graphics* for drawing techniques. See Section 12.6.3 in Lesson 12 for horizontal grouping signs.

For further practice, see Addendum 1—Reading Practice.

Submit Exercise 17 to your instructor.
--

BLANK PAGE

PRACTICE 17D

1
$$2x^2 + 3x - 5 = x^2 + 7x + 10$$

2

3
$$2x^2 + 3x - 5 = x^2 + 7x + 10$$

4
$$2x^2 + 3x - 5 = x^2 + 7x + 10$$

5

6
$$2x^2 + 3x - 5 = x^2 + 7x + 10$$

7
$$2x^2 + 3x - 5 = x^2 + 7x + 10$$

8
$$2x^2 + 3x - 5 = x^2 + 7x + 10$$

9

10
$$2x^2 + 3x - 5 = x^2 + 7x + 10$$

11
$$2x^2 + 3x - 5 = x^2 + 7x + 10$$

12

13
$$2x^2 + 3x - 5 = x^2 + 7x + 10$$

14

15
$$2x^2 + 3x - 5 = x^2 + 7x + 10$$

16
$$2x^2 + 3x - 5 = x^2 + 7x + 10$$

17
$$2x^2 + 3x - 5 = x^2 + 7x + 10$$


18
$$2x^2 + 3x - 5 = x^2 + 7x + 10$$


19


20
$$2x^2 + 3x - 5 = x^2 + 7x + 10$$


Line 4: A space is inserted before x and y because terms are aligned, as printed.
Lines 6-7: Although the equals signs are aligned in print, terms are not aligned. In braille, each line is left adjusted.
Line 8: An additional transcriber-inserted grouping symbol is needed because the remark takes three lines.
Lines 15-18: Although this is not a system of equations (there is only one variable), alignment guidelines in [Section 17.2\(c\)](#) are followed. Terms are aligned on the left of the equals signs; to the right, place values are not aligned.


PRACTICE 17I


1 


2 


3 


4 


5 


6 


7 


8 


9 


10 

11 

12 












13 

14 

15 

Notice the placement of the blank lines. Blank lines 2 and 6 precede and follow the spatial material on lines 3-5; blank lines 10 and 14 precede and follow the spatial material on lines 11-13. The code switch indicators are placed on either side of those blank lines.

PRACTICE 17K

1 
2 
3 
4 
5 
6 
7 
8 
9 
10 
11 
12 